

## 6His, His-Tag Monoclonal antibody

**Cat:**B30001T

**Company:** HaoKebio

**Uniprot ID:**HISTAG

**Applications:** IF/ICC : 1:100-1:400

**Organism:**Mouse


WB:1:2500-1:25000

**Species reactivity:**Recombinant protein

IP : 1ug for 1mg of total protein lysate

**Background:**

Protein tags are a protein or peptide sequences located either on the C- or N- terminal of the target protein. His-tag is often used for affinity purification and binding assays. Expressed. The His-tag antibody is a useful tool for monitoring of the His-tagged proteins and recognizes His-tags placed at N-terminal, C-terminal, and internal regions of fusion proteins expressed in bacteria, insect, and mammalian cells. A His-tag (polyhistidine tag) consists of at least six histidine residues that are located at the N- or C-terminus of recombinant proteins. It is commonly used for affinity purification and protein binding experiments.

**Images:**


Dilution of 1:25000 incubated at room temperature for 1.5 hours.

**Source of Reagents:**

发表[中文论文]请标注:6His, His-Tag(B30001T)由杭州浩克生物技术有限公司提供;

发表[英文论文]请标注:6His, His-Tag(B30001T) were kindly provided by Hangzhou Haoke Biotechnology Co., Ltd.

**Protein full name:**

6\*His, His-Tag

**Synonyms:**

6\*His, His Tag, His Tag, 6\*His, 6\*his tag, 6\*his, His tag

**Immunogen:**

Peptide

**Isotype:**

IgG1

**Purity:**

Protein G purification

**Form:**

Liquid

**Storage Buffer:**

PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.

**Storage:**

Store at -20 °C for one year.