

HRP-conjugated GAPDH Monoclonal antibody

Cat:B31013T-HRP

Company: HaoKebio

Uniprot ID:P04406

Applications:

WB:1:2500-1:25000

Organism:Mouse

Species reactivity:Human Mouse Rat

Molecular Weight Calculation: 36 kDa

Observed Molecular Weight: 36 kDa

Background:

Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) catalyzes the phosphorylation of glyceraldehyde-3-phosphate during glycolysis. GAPDH participates in nuclear events including transcription, binding RNA, RNA transportation, DNA replication, DNA repair and apoptosis. Being stably and constitutively expressed at high levels in most tissues and cells, GAPDH is considered a housekeeping protein. It is widely used as a control for RT-PCR and also loading control in electrophoresis and Western blotting. GAPDH is normally expressed in cellular cytoplasm or membrane, but can occasionally translocate to the nucleus after the addition of post-translational modifications such as S-nitrosylation. This antibody is raised against full length GAPDH of human origin. It can recognize the 36 kDa GAPDH protein in most cells/tissues. In addition, a band below 36 kDa can always be detected as the isoform or spliced product of GAPDH. Please note that some physiologic factors, such as hypoxia and diabetes, increase GAPDH expression in certain cell types.

Affinity purification
Form:


Liquid

Storage Buffer:

PBS with 0.02% sodium azide, 100 µg/ml BSA and 50% glycerol.

Storage:

Store at -20 °C for one year.

Images:


Dilution of 1:25000 incubated at room temperature for 1.5 hours.

Source of Reagents:

发表[中文论文]请标注:HRP-GAPDH (B31013T-HRP)由杭州浩克生物技术有限公司提供;

发表[英文论文]请标注:HRP-GAPDH (B31013T-HRP) were kindly provided by Hangzhou Haoke Biotechnology Co., Ltd.

Protein full name:

glyceraldehyde-3-phosphate dehydrogenase

Synonyms:

1H18

Immunogen:

Recombinant protein

Isotype:

IgG2b

Purity: