

Proteasome 26S S10 Recombinant antibody

Cat:B35098D

Company: HaoKebio

Uniprot ID:O75832

Applications: IHC:1:50-1:100

Organism:Rabbit

IHC-Polymer:1:200-1:400

Species reactivity:Human Mouse Rat

IHC-TSA:1:250-1:500

Molecular Weight Calculation: 24 kDa


WB:1:1000-5000

Observed Molecular Weight: 24 kDa

Background:

This gene encodes a subunit of the PA700/19S complex, which is the regulatory component of the 26S proteasome. The 26S proteasome complex is required for ubiquitin-dependent protein degradation. This protein is a non-ATPase subunit that may be involved in protein-protein interactions. Aberrant expression of this gene may play a role in tumorigenesis. Two transcripts encoding different isoforms have been described. Pseudogenes have been identified on chromosomes 3 and 20.

dium high heat for 8 minutes. Incubate antibody, 4°C overnight. Secondary antibody: Poly-HRP Goat Anti-Rabbit & Mouse Universal Secondary Antibody, RT, 1h.

Images:

Synonyms:

p28; p28(GANK); dj889N15.2

Dilution of 1:5000 incubated at room temperature for 1.5 hours.

Immunogen:

Recombinant protein

Source of Reagents:

发表[中文论文]请标注:Proteasome 26S S10(B35098D)由杭州浩克生物技术有限公司提供;

发表[英文论文]请标注:Proteasome 26S S10(B35098D) were kindly provided by Hangzhou Haoke Biotechnology Co., Ltd.

Isotype:

IgG

Subcellular location:

Cytoplasm,Nucleus

Purity:

Affinity purification

Form:

Liquid

Storage Buffer:

PBS with 0.02% sodium azide, 100 µg/ml BSA and 50% glycerol.

Storage:

Store at -20 °C for one year.

Experimental procedure:

Antigen retrieval: Citrate buffer (pH 9.0) , Medium high heat for 8 minutes, stop for 7 minutes, me