

VCP Recombinant antibody

Cat:B36240S
Company: HaoKebio

Uniprot ID:P55072

Applications: IHC:1:50-1:500

Organism:Rabbit

IHC-Polymer:1:200-1:2000

Species reactivity:Human Mouse Rat

IHC-TSA:1:250-1:2500

Molecular Weight Calculation: 89 kDa

WB:1:5000-1:50000

Observed Molecular Weight: 89 kDa

Background:

VCP(Valosin-containing protein), also known as TER ATPase and 15S Mg²⁺-ATPase p97 subunit, belongs to the AAA ATPase family. VCP was first identified as a result of attempts to clone a putative peptide hormone called valosin. It was found that the cloned cDNA encoded a ubiquitously expressed 90 kDa cytosolic protein, termed VCP, which showed none of the characteristics of a peptide hormone precursor. Defects in VCP are the cause of inclusion body myopathy with early-onset Paget disease and frontotemporal dementia (IBMPFD) and amyotrophic lateral sclerosis type 14 with or without frontotemporal dementia (ALS14). VCP has a calculated molecular weight of 89 kDa and an apparent molecular weight of 90-100 kDa.

PBS with 0.02% sodium azide,100 µg/ml BSA and 50% glycerol.

Storage:


Store at -20 °C for one year.

Experimental procedure:

Antigen retrieval: Citrate buffer (pH 9.0) , Medium high heat for 8 minutes, stop for 7 minutes, medium high heat for 8 minutes. Incubate antibody, 4°C overnight. Secondary antibody: Poly-HRP Goat Anti-Rabbit & Mouse Universal Secondary Antibody, RT, 1h.

Images:


Sample: Mouse testis, 4% PFA 12-24h



Dilution of 1:10000 incubated at room temperature for 1.5 hours.

Synonyms:

HEL-S-70, HEL-220, EC:3.6.4.6, 3G23, 15S Mg²⁺-ATPase p97 subunit

Immunogen:

Recombinant protein

Isotype:

IgG

Subcellular location:

Cytoplasm,Nucleus

Purity:

Affinity purification

Form:

Liquid

Storage Buffer:

Source of Reagents:

发表[中文论文]请标注:VCP(B36240S)由杭州

浩克生物技术有限公司提供;

发表[英文论文]请标注:VCP(B36240S) were kindly provided by Hangzhou Haoke Biotechnology Co., Ltd.